

Secompla 67C Binding Polymer

technology sheet

Produce superior adhesion of seed treatments and colourants to canola with Secompla™ 67C binding polymer.

- High-quality product for uniform colour and treatment coverage
- Control dust-off during treatment, processing, planting and re-bagging with low application rates
- Increase seed protection by binding the colour and seed treatment onto the seed for a professional appearance

Active ingredient

Secompla 67C

Formulation

Water-based liquid concentrate with neutral pH

One case contains

2 x 9.46 L jug
Also available in 984.2 L tote

Storage

2-year shelf life. Store away from heat. Protect from direct sunlight and freezing.

Crops

Canola

Treatment

Applied directly on seed

How it works

Secompla 67C binding polymer provides an improvement to the adhesion of the active ingredient(s) on seed, along with any other components, such as seed colourants. By providing greater binding strength, this ensures that seed treatments stay where they belong for maximum performance.

This additional binding strength also assists in reducing dust-off during treatment, processing, planting and re-bagging of canola seed, while ensuring a professional look to canola seed.

Some variation may occur. Adhesion of chemical seed treatments and colourants on seed will differ based on the following considerations:

- application rate
- seed shape
- chemical seed treatments included
- seed size
- crop type
- seed texture



Untreated



Secompla 67C Binding Polymer

Source: BASF, 2017.

 **BASF**
We create chemistry

Application rates

One case treats 24,194 (24.2 tonnes) to 145,538 kg (145.5 tonnes) of seed. One tote treats 1,258,568 (1,258.6 tonnes) to 7,570,769 kg (7,570.8 tonnes) of seed.

Crop	ml per 100 kg of Seed		kg of Seed Treated Per Package Size by Application Rate			
	Low Rate	High Rate	2 x 9.46 L		1 x 984.2 L	
			Low Rate	High Rate	Low Rate	High Rate
Canola (treated)	32.5	65.0	58,215	29,108	3,028,308	1,514,154
Canola (5% wt. grain)	13.0	78.2	145,538	24,194	7,570,769	1,258,568

Mixing order

Slurry Mix Method

1. Shake well or agitate prior to use.
2. Start and maintain constant agitation throughout entire mixing and treatment process.
3. Add seed treatment materials per their label instructions.
4. Add Color Coat colourant and Secompla 67C.
5. Fill water volume to desired volume if required.
6. Mix treatment solution for 10 minutes following the addition of the colourant.

Application tips

Begin at low rate and evaluate if more product is needed to obtain desired colour.

Do not apply to food or feed crops or grain after harvest. A spray shield is recommended for applications if over-spray or drift is a potential problem.

Do not discharge into subsoil/soil, drains, surface water and groundwater.

Tank mixes

Always check compatibility of all components in the treatment slurry prior to large scale mixing.

Contact **AgSolutions**® Customer Care at 1-877-371-BASF (2273) or your local BASF Sales Representative for additional information.

Always read and follow label directions.