

Sercadis fungicide

technology sheet

grapes

Innovative, new chemistry for consistent, continuous control of powdery mildew.

- Control of powdery mildew in grapes
- Tank-mix and rotational flexibility to adapt to the season's needs
- Highly systemic activity helps protect new growth

Active ingredient

Fluxapyroxad – Group 7

Formulation

Suspension

One case contains

2 x 1.35 L jugs

Storage

Store away from food or feed.

Crop

Grapes

Timing

Begin prior to onset of disease
on a 7 to 14 day interval

Disease controlled

In grapes.

Powdery mildew (*Erysiphe necator*)



Application rates

One jug of Sercadis® fungicide treats 5.4 ha (13.5 acres).

Grapes

Powdery mildew 250 ml/ha (100 ml/ac)

Mixing order

1. Ensure the spray tank is clean before use.
 2. Fill the tank 1/2 full of water and start agitation.
 3. Add the required amount of Sercadis fungicide to the tank.
 4. Add the required amount of tank mix partner, if applicable.
 5. Continue agitation while filling the remainder of the spray tank.
 6. After use, clean the spray tank according to label precautions.
-

Application tips

Rainfastness – 1 hour.

Restricted entry interval – 12 hours.

Resistance management – Tank-mix with a non-Group 7 fungicide when such use is permitted. Avoid consecutive applications of Sercadis or other Group 7 fungicides in a season.

Pre-harvest interval

14 days for grapes.

Tank mixes

None on label.

Contact **AgSolutions®** Customer Care or your BASF Sales Representative for information on supported tank mixes.

For more information

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca.

Always read and follow label directions.

AgSolutions is a registered trade-mark of BASF Corporation; SERCADIS is a registered trade-mark of BASF SE; both used with permission by BASF Canada Inc. SERCADIS fungicide should be used in a preventative disease control program. © 2016 BASF Canada Inc.