



We create chemistry

# Priaxor<sup>®</sup>

## Xemium<sup>®</sup> Fungicide

### BASF Contact

Sydney Marlow  
Canola Crop Manager  
BASF Canada Inc.  
905-301-0073  
[sydney.marlow@basf.com](mailto:sydney.marlow@basf.com)

### What is PRIAXOR?

- A Xemium fungicide that provides consistent, continuous disease control.

### Active Ingredients

- Xemium—Group 7
- Pyraclostrobin—Group 11

### Registered Crops

- |  |  |
|--|--|
| <ul style="list-style-type: none"><li>• Canola</li><li>• Soybean</li><li>• Corn</li><li>• Flax</li><li>• Alfalfa (seed production)</li><li>• Grasses (seed production)</li><li>• Sunflower</li><li>• Wheat</li></ul> | <ul style="list-style-type: none"><li>• Barley</li><li>• Rye</li><li>• Oats</li><li>• Chickpea</li><li>• Dry bean</li><li>• Faba bean</li><li>• Field Pea</li><li>• Lentil</li></ul> |
|--|--|

### Diseases Controlled

- |  |  |
|--|--|
| <ul style="list-style-type: none"><li>• Blackleg</li><li>• White mold (suppression)</li><li>• Asian soybean rust</li><li>• Frogeye leaf spot (suppression)</li><li>• <i>Septoria</i> brown spot</li><li>• Common rust</li><li>• Gray leaf spot</li><li>• Northern leaf blight</li><li>• <i>Alternaria</i> black spot</li></ul> | <ul style="list-style-type: none"><li>• Common leaf spot</li><li>• Blossom blight (suppression)</li><li>• Pasmó</li><li>• <i>Sclerotinia</i> stem rot (suppression)</li><li>• Leaf and stem rust</li><li>• Powdery mildew</li><li>• <i>Anthraco</i></li><li>• <i>Mycosphaerella</i> blight</li><li>• <i>Ascochyta</i> blight</li></ul> |
|--|--|



We create chemistry

### Application Timing

- |  |   |
|--|---|
| <ul style="list-style-type: none"><li>• Soybean: early to late flower (R2 to R3)</li><li>• Corn: early to full tassel (VT)</li><li>• Canola:<ul style="list-style-type: none"><li>○ 2 to 6 leaf (rosette) stage for Blackleg control</li><li>○ 20 to 50 per cent bloom for <i>Alternaria</i> black spot and <i>Sclerotinia</i> stem rot suppression</li></ul></li><li>• Flax: 20 to 50 per cent flowering</li><li>• Alfalfa: 10 to 30 per cent bloom</li></ul> | <ul style="list-style-type: none"><li>• Grasses: prior to onset of symptoms</li><li>• Sunflowers: at onset of symptoms</li><li>• Chickpeas: beginning of flowering</li><li>• Dry beans: beginning of flowering or at onset of symptoms</li><li>• Field peas: beginning of flowering or at onset of symptoms</li><li>• Lentils: beginning of flowering or at the onset of symptoms</li></ul> |
|--|---|

### Feature Benefits

- Crops treated with PRIAXOR experience better management of minor stresses.
- Includes proven benefits of AgCelence, which often result in plants that are larger and greener with thicker stems, larger root mass, and increased yield potential.
- Xemium is active on key diseases and has unique mobility characteristics. Because of this, it can better penetrate the various levels within the leaf structure. It also forms deposits on the leaves for later movement and disease control.